

Lower St. Louis River Habitat Plan

Actions:

- The Great Lakes Panel on Aquatic Nuisance Species should continue work to strengthen federal legislation through reauthorization of NISA.
- The International Joint Commission, Coast Guard, and shipping industry should develop a comprehensive ballast water control plan. The plan should be coordinated throughout the Great Lakes states to ensure a consistent policy.
- Develop new technologies to treat and/or manage ballast water.

Strategy 15: Develop Zebra Mussel Control Strategy

Problem Summary: Introduced via ballast water in the late 1980s, the population of the non-native zebra mussel is currently expanding in the Lower St. Louis River. Zebra mussels attach themselves in dense layers to any hard surface, including native mussels, aquatic plants, surface water intake pipes, docks, and pilings. In addition to competing with native mussels for food, the zebra mussels cover native mussels so densely that they are unable to open and close their shells. Left unchecked, zebra mussels are likely to eliminate populations of native mussels throughout the harbor, and possibly upstream to the Fond du Lac dam.

Strategy: Develop a zebra mussel control strategy that includes preventing upstream expansion of zebra mussel populations and controlling or eliminating existing populations of zebra mussels.

Actions:

- Complete survey field work to determine habitat needs of native mussels.
- Identify important native mussel beds, map them, and eliminate zebra mussels.
- Establish a long-term monitoring protocol for zebra and native mussels; monitor the presence, locations, and abundance of native and non-native mussel species; determine upstream limits to distribution of zebra mussels, and determine the influence of seiche in distribution of zebra mussels.
- Determine water flow in the estuary and make predictions/assessments about whether zebra mussels will colonize and infest the entire Lower St. Louis River.
- Consult with experts from other areas, such as Lake Erie, the Mississippi River, and the St. Croix River, to help assess the likelihood of zebra mussels spreading in the Lower St. Louis River.

Strategies to Address Multiple Stresses

Strategy 16: Infuse the Lower St. Louis River Habitat Plan into Public Planning Processes

Problem Summary: From an ecological perspective, most of the major stresses to the Lower St. Louis River—loss of habitat, increased sedimentation, degradation of water quality, and exposure to sediment-associated contaminants—are directly related to land use decisions, but land use decisions are often based on incomplete or inaccurate ecological information. Rarely does the public planning process include a comprehensive determination of the effects that proposed land uses will have on the overall health of the watershed.

Strategy: The Lower St. Louis River Habitat Plan should be infused into public planning processes throughout the region. The focus should be on those plans that are directly linked to the estuary and the immediate surrounding communities of Superior and Duluth, such as the Port Plan and city comprehensive land use plans. Science-based information needs to be an integral part of the public planning process throughout the entire St. Louis River watershed.

Actions:

- Identify local planning initiatives within the Lower St. Louis River watershed, including but not limited to, the Port Plan, Port Management Plan, city comprehensive plans, and management plans for recreation areas.
- Once identified, members of the Habitat Plan partnership should be matched with each local planning initiative.

Strategy 17: Encourage the Development and Implementation of a Comprehensive Port Plan

Problem Summary: Commercial shipping and associated industries are a vital part of the Twin Ports economy. However, development projects, dredging and filling, and other commercial activities can result in the loss and degradation of critical estuarine aquatic habitats that support fish, mussels, and breeding and migratory birds. Balancing effective utilization of the waters, near shore areas, and waterfront lands of the Duluth-Superior Harbor while also protecting, restoring, and enhancing the valuable habitat of the Lower St. Louis River is a difficult task.

Strategy: A comprehensive Port Plan should be developed and implemented. A carefully developed and broadly accepted comprehensive Port Plan can help to provide needed balance. An effective Port Plan will provide the basis for protection, restoration, and enhancement of the many assets of the Duluth-Superior Harbor and will help assure that commerce and recreation can develop fully without impairment of the habitats that are so vital to the health of the western Lake Superior region. An effective Port Plan will identify sensitive areas where extra care is needed to preserve, restore, and enhance the values of the Lower St. Louis River, including critical and important habitat. An effective Port Plan will result in preservation of waterfront areas for uses that are consistent with the existing infrastructure, including the navigation channels and associated structures. It is very important that maritime

Lower St. Louis River Habitat Plan

commerce utilize existing harbor infrastructure rather than encroach on the remaining high quality habitat areas of the Lower St. Louis River. Deep-water maritime commerce should be consolidated along the existing 28-foot channels and should use existing structures as much as possible. Land fronting the existing 23-foot channels should be reserved for maritime commerce that can utilize these channels without the requirement for deepening. Expansion of constructed and maintained navigation facilities should be carefully evaluated in light of the habitat values inherent in the Lower St. Louis River. The remaining waterfront land should be reserved for water-dependent activities including habitat protection, recreation, and scenic beauty. Land uses that are not water dependent should be restricted to non-waterfront sites. Plans should be developed to protect sensitive areas from spills and other accidents or failures. These plans should be included in the Port Plan directly or by reference.

Actions:

- Support the development of an effective Port Land-use Plan for the Superior, Wisconsin, portion of the Harbor. Complete review and acceptance of the Superior Port Plan will be important so that it can be used as a template for updating the Duluth Port Plan.
- Support revision of the Duluth Port Plan so the two plans form a comprehensive guide to effective management and utilization of the values of the Lower St. Louis River and the Duluth-Superior Harbor.

Strategy 18: Encourage the Development and Implementation of a Comprehensive Dredge Material Management Plan

Problem Summary: A century of dredging and disposing of dredged materials has extensively modified habitat in the Lower St. Louis River. Ongoing dredging and dredged materials management practices have the potential to cause additional loss and degradation of estuarine aquatic habitats and place additional stress on fish, mussel, and bird populations.

Strategy: A comprehensive dredged materials management plan (DMMP) should be developed and implemented. A draft DMMP was developed in 1997. This draft should be updated and approved. A good comprehensive DMMP can result in reduced impacts to habitats from dredging practices. While restoration of the habitat in the Lower St. Louis River to pre-dredging conditions is not feasible and is not a goal of this Habitat Plan, future dredged materials management activities should be designed and managed to minimize detrimental impacts to important habitats. Opportunities to protect, restore, or enhance habitat through the control and redesign of dredged material management practices should be fully exploited.

An effective dredged materials management plan will

- enable habitat restoration and enhancement that may be economically difficult or impossible without the benefit of redirected dredged materials management activities;
- reduce the need for dredging by recommending land use changes that will result in reduced sedimentation within the harbor;
- provide the basis to obtain additional funding to demonstrate and practice innovative beneficial reuse of dredged materials and important habitat restoration and enhancement activities.

Lower St. Louis River Habitat Plan

Beneficial reuse of dredged materials to restore and enhance habitat may facilitate regulatory approval of dredging projects that would be difficult or impossible to authorize if containment or disposal of the dredged materials was proposed. It is also possible that streamlined regulatory processes can be utilized for projects that are consistent with an agency-approved comprehensive dredged materials management plan. Therefore, a comprehensive dredged materials management plan will benefit commercial interests in the harbor in addition to natural resources interests in the Lower St. Louis River.

Actions:

- A comprehensive dredged materials management plan should be developed cooperatively by the Port Authority, Corps of Engineers, WDNR, MDNR, MPCA, the CAC, and others. The Harbor Technical Advisory Committee (HTAC) of the Metropolitan Interstate Committee (MIC) will provide the best forum to bring the necessary partners together for this purpose. Considerable work has already been produced that can provide the basis for an effective dredged materials management plan.



Lower St. Louis River Habitat Plan



INDICATORS OF SUCCESS

Once the implementation of the Habitat Plan is underway, it is critical to periodically assess whether the strategies being implemented are successfully mitigating threats and improving the health of the conservation targets. If the strategies implemented are not meeting the conservation goals of each target—typically, improving or maintaining the health of species and ecosystems—it is necessary to revise strategies. Monitoring the responses to management strategies and adjusting the strategies accordingly is referred to as “adaptive management.” In this Habitat Plan, “adaptive management” includes adjustments not only to traditional ecosystem management strategies (e.g., prescribed fires, wetland restoration), but to the entire suite of strategies developed through this planning effort.

There are three levels at which the success of implementation of the Habitat Plan should be measured:

- health of the conservation targets;
- mitigation of the threats to the conservation targets; and
- implementation of the strategies.

Change in the health of conservation targets (plant communities, aquatic habitats, species) is the most fundamental measure of the successful implementation of any conservation plan. However, measurable changes for many of the conservation targets in this Plan will occur over widely varying time periods. For many conservation targets of the Lower St. Louis River, changes will not be measurable within three to five years of implementing the strategies intended to improve their health. At one extreme, the initial changes to upland forests may not be noticeable for a decade or more, and significant, long-term changes may not be measurable for several decades. At the other extreme, initial changes to piping plover populations could be noticeable within the first three to five years of implementing strategies intended to improve their health.

Since it will take some time for the health of conservation targets to show improvement, a second important indicator to monitor is how well threats to the conservation targets are mitigated. The mitigation of threats will be measurable somewhat earlier than changes in the overall health of conservation targets.

Finally, it is necessary to track the strategies outlined in this Plan to ensure that the appropriate actions are taken to successfully implement them. Since more than one strategy may improve the health of a single conservation target, it may be not be immediately clear which strategy is responsible for improvements in the health of that conservation target. Monitoring at all three levels—health of conservation targets, mitigation of threats, and implementation of strategies—will help clarify which strategies are successful and which strategies need to be revised or eliminated. Although the ultimate effects of each strategy will take several years to be measurable, the implementation of strategies should be tracked immediately.

To effectively assess changes in the health of conservation targets, it is critical for each monitoring effort to be clearly linked to the factors that define the health of the conservation targets. Those factors are described in the earlier text on the conservation targets and are also summarized in Table 6. A preliminary list of broadly defined monitoring efforts is included. Though extensive, this is not a final or comprehensive list of all necessary monitoring efforts. It represents priority areas where work should be initiated. Beginning to monitor the factors identified will be necessary to both measure success and refine future monitoring efforts.

Lower St. Louis River Habitat Plan

Table 6. Factors indicating the health of the conservation targets and preliminary considerations for monitoring those factors.

*This is not a comprehensive list.

Conservation Target	Indicator of Health to be Measured	Preliminary Monitoring Considerations*
<p>Large riverine reach</p>	<p>Hydrologic regime (volume, rate, timing of river flow) Water quality (sediments, nutrients, chemistry) Habitat morphology Diversity and relative abundance of fish species Diversity and relative abundance of bird species Lake sturgeon spawning, reproduction</p>	<p>Develop hydrologic model for estuary that includes regime under natural range of variation. Sample peak, low, and base flows on periodic basis; compile seasonal and annual measurements and compare to hydrologic model under natural range of variation. Develop model of sediment and nutrient loads to estuary; describe loads expected under natural range of variation. Measure sediment and nutrient loads under range of conditions (“normal,” post-storm, during spring thaw). Sample water chemistry. Periodically assess seasonal fish use and abundance in habitat. Periodically assess seasonal bird use and abundance in habitat. See also lake sturgeon.</p>
<p>Upper estuarine (undredged) river channel</p>	<p>Hydrologic regime (volume, rate, timing of river flow) Water quality (sediments, nutrients, chemistry) Habitat morphology Diversity and relative abundance of fish species Diversity and relative abundance of bird species</p>	<p>See large riverine reach.</p>
<p>Lower estuarine (dredged) river channel Lower estuary (industrial harbor) flats</p>	<p>Hydrologic regime (volume, rate, timing of river flow) Water quality (sediments, nutrients, chemistry) Diversity and relative abundance of fish species Diversity and relative abundance of bird species</p>	<p>See large riverine reach.</p>

Lower St. Louis River Habitat Plan

<p>Upper estuary flats Sheltered bays Clay-influenced river mouths Clay-influenced bay</p>	<p>Hydrologic regime (volume, rate, timing of river flow) Water quality (sediments, nutrients, chemistry) Habitat morphology Diversity and relative abundance of fish species Diversity and relative abundance of bird species Diversity and relative abundance of wetland plant species Extent (acreage) of wetland vegetation</p>	<p>See large riverine reach. Conduct field surveys every three years; record native species composition, presence and abundance of any non-native species, and overall health ranking. Photo-monitoring should be part of these field surveys. Air photo interpretation every five years should be used to assess patterns of various wetland types (e.g., emergent marsh, submergent marsh, wet meadow, etc.) and compare changes in extent of the various wetland plant communities within the estuarine aquatic habitats.</p>
<p>Industrially-influenced bays</p>	<p>Hydrologic regime (volume, rate, timing of river flow) Water quality (sediments, nutrients, chemistry) Habitat morphology Diversity and relative abundance of fish species Diversity and relative abundance of bird species Diversity and relative abundance of wetland plant species Extent (acreage) of wetland vegetation (restored bays or slips)</p>	<p>See upper estuary flats, sheltered bays, clay-influenced river mouths, and clay-influenced bay. May not apply to all industrially-influenced bays.</p>
<p>Industrial slips</p>	<p>Water quality (sediments, nutrients, chemistry) Habitat morphology Diversity and relative abundance of fish species Diversity and relative abundance of bird species</p>	<p>Sample water chemistry. Periodically assess seasonal fish use and abundance in habitat. Periodically assess seasonal bird use and abundance in habitat.</p>
<p>Clay-influenced tributaries Bedrock-influenced tributaries</p>	<p>Hydrologic regime (volume, rate, timing of river flow) Water quality (sediments, nutrients, chemistry) Habitat morphology Diversity and relative abundance of fish species</p>	<p>See large riverine reach.</p>

Lower St. Louis River Habitat Plan

Estuarine Plant Community Targets		
Great Lakes coastal wetland complex	Diversity and relative abundance of wetland plant species Rare plant species Lack of non-native species Extent (acreage) of wetland vegetation Spatial patterns of wetland plant community types (e.g., emergent marsh, submergent marsh, wet meadow, etc.)	See upper estuary flats, sheltered bays, clay-influenced river mouths, and clay-influenced bay.
Baymouth Bar Community Targets		
Beaches	Erosion and deposition of sand	Air photo or satellite imagery interpretation every five years to assess erosion and deposition patterns on Points and around western Lake Superior coastline.
Beachgrass dunes Dune shrublands Interdunal wetlands	Diversity and relative abundance of plant species Rare plant species Extent (acreage) of each community type Lack of non-native species	Conduct field surveys every three years; record native species composition, presence and abundance of any non-native species, and overall health ranking. Photo-monitoring should be part of these field surveys. Air photo interpretation every ten years should be used to assess extent and spatial patterns of these community types.
Dune pine forests	Diversity and relative abundance of plant species Rare plant species Extent (acreage) of each community type (increased from current area) Lack of non-native species	Conduct field surveys every three years; record native species composition, presence and abundance of any non-native species, and overall health ranking. Photo-monitoring should be part of these field surveys. Air photo interpretation every ten years should be used to assess extent and spatial patterns of this community type.
Upland Forest Community Targets		
White pine-red pine forests Northern conifer-hardwoods forest / Northern hardwoods forest Spruce-fir boreal forest	Diversity and relative abundance of plant species Age class structure of dominant tree species Extent (acreage) of each community type (increased from current area)	Map land cover and complete a change detection analysis, using satellite imagery, to assess species composition. Conduct field surveys to evaluate age class structure; every ten years may be an appropriate frequency.

Lower St. Louis River Habitat Plan

Other Inland Plant Community Targets		
Eroding clay bluffs Clay seeps Conifer swamps Hardwood swamps Shrub swamps Inland marshes Wet meadows Fens	Diversity and relative abundance of plant species Hydrologic regime (volume, rate, timing, source of flow)	Conduct field surveys to evaluate the species diversity and structure of these finer-scale communities. Photo monitoring may be a good technique for assessing the eroding clay bluffs and clay seeps.
Cliffs and rock outcrops	Diversity and relative abundance of plant species	Conduct field surveys to evaluate the species diversity and structure of these finer-scale communities.
Species Targets		
Native fish assemblage	Diversity and relative abundance of native fish species Non-native aquatic animal species declining or eradicated	Continue existing fisheries sampling efforts; determine whether additional, more comprehensive sampling is necessary.
Lake sturgeon	Population size, age class structure	Continue WDNR's and MDNR's monitoring of population size and reproductive success; expand to include radio-tag monitoring.
Native mussel assemblage	Diversity and relative abundance of native mussel species Zebra mussel controlled or eradicated	Continue MDNR's mussel sampling effort to develop estimates of current populations of native species; continue field sampling to monitor both native and zebra mussel populations.
Migratory bird aggregations Breeding bird aggregations	Diversity and numbers of migratory birds Diversity and numbers of breeding birds	Coordinate with existing survey and other sampling efforts. Conduct a comprehensive breeding survey to estimate current breeding status and populations of breeding birds. Conduct periodic monitoring to estimate long-term trends. Conduct a comprehensive migratory survey to estimate current diversity and numbers of birds utilizing estuary.
Piping plover	Population size, reproduction	Conduct detailed annual surveys; coordinate with U.S. FWS recovery and monitoring efforts.
Common tern	Population size, reproduction	Conduct detailed annual surveys; coordinate with U.S. FWS recovery and monitoring efforts.
Wild rice	Population size/areal extent, reproduction	Conduct field surveys.

*This is not a comprehensive list.

Lower St. Louis River Habitat Plan

Effective evaluations of how well threats are being mitigated also requires a clear link between the threats and the factors that indicate mitigation. These factors are summarized in Table 7.

Table 7. Threats and factors to be evaluated that indicate how well threats are being mitigated.

Stress	Sources of Stress	Indicator of Mitigation to be Measured
Loss of habitat	Development Commercial shipping (dredging and filling) Other sources	# of acres developed in/on “natural” communities or habitats (vs. acres redeveloped within existing developed areas). Monitor whether dredged area expands or remains stable.
Increased sedimentation	Development Forest management practices Other sources	Change in sediment load.
Competition from non-native species	Commercial shipping Development (accidental release or dispersal of non-native species) Other sources	Rate of introduction of new non-native species declines, or new introductions are eliminated. Populations of non-native species decline, are eliminated, or are controlled.
Exposure to sediment-associated contaminants	Contaminated sediments (from historical, municipal sewage, commercial, and industrial releases) Other sources	Acreage of highly contaminated sediments (acreage should decrease as problem is addressed).
Degraded water quality	Development Commercial shipping Contaminated sediments (from historical, municipal sewage, commercial, and industrial releases) Forest management practices Other sources	Water quality measures improve from current levels.

Measuring the health of conservation targets and the level of threat mitigation requires integration of monitoring activities and data across agencies and jurisdictional borders. Implementation of an ecological monitoring program should include application of a strong, statistically robust design and should utilize Geographic Information Systems (GIS) for analysis and display of results. Ideally, all of the specific examples described above will be part of a single coordinated monitoring effort among many agencies. It will be necessary for this integrated program to go beyond the levels of monitoring and coordination currently being done in the Lower St. Louis River.

REFERENCES

- Allen, James. 1832. The journal of Lieutenant James Allen, expedition of 1832. U.S. House Executive Documents, No. 323, 23^d Cong., 1 Sess, pp. 7-68.
- ASCI Corporation. 1999. Summary of test results determining potential mercury, PAH and PCB bioaccumulation by *Lumbriculus variegatus* exposed to St. Louis Bay sediment samples. ASCI Corporation, Duluth, MN.
- Baker, F. C. 1918. The relation of shellfish to fish in Oneida Lake, New York. Syracuse: New York State College of Forestry at Syracuse University, Circular (21):11-34.
- Blasland Bouck and Lee, Inc. 2000. Supplemental surface water and streambed sediment investigation report for Koppers Industries, Inc. facility, Superior, WI.
- Breneman, D., C. Richards, and S. Lozano. 2000. Environmental influences on benthic community structure in a Great Lakes embayment. *J. Great Lakes Res.* 26(3):287-304.
- Bronte, C.R., L.M. Evrard, W.P. Brown, K.R. Mayo, and A.J. Edwards. 1998. Fish community changes in the St. Louis River estuary, Lake Superior, 1989-1996: Is it ruffe or population dynamics? *J. Great Lakes Res.* 24(2):309-318.
- Canham, C.D., and O.L. Loucks. 1984. Catastrophic windthrow in the presettlement forests of Wisconsin. *Ecology* 65(3):803-809.
- Carroll, L.M., and F.R. Raiter. 1990. The fires of autumn: The Cloquet-Moose Lake disaster of 1918. St. Paul: Minnesota Historical Society Press.
- Chotkowski, M.A., and J.E. Marsden. 1999. Round goby and mottled sculpin predation on lake trout eggs and fry: Field predictions from laboratory experiments. *J. Great Lakes Res.* 25(1): 26-35.
- Crane, J.L. 1999. Assessment of contaminated sediments in Slip C, Duluth Harbor, Minnesota. EPA-905-R-99-007. U.S. Environmental Protection Agency, Great Lakes National Program Office, Chicago, IL.
- Crane, J.L. 2001. Quality assurance project plan (QAPP): GIS-based contaminated sediment database for the St. Louis River Area of Concern. Environmental Outcomes Division, Minnesota Pollution Control Agency, St. Paul, MN.
- Crane, J.L., M. Schubauer-Berigan, and K. Schmude. 1997. Sediment assessment of hotspot areas in the Duluth/Superior Harbor. EPA-905-R97-020. U.S. Environmental Protection Agency, Great Lakes National Program Office, Chicago, IL.

Lower St. Louis River Habitat Plan

- Crane, J.L., D.D. MacDonald, C.G. Ingersoll, D.E. Smorong, R.A. Lindskoog, C.G. Severn, T.A. Berger, and L.J. Field. 2000. Development of a framework for evaluating numerical sediment quality targets and sediment contamination in the St. Louis River Area of Concern. EPA 905-R-00-008. U.S. Environmental Protection Agency, Great Lakes National Program Office, Chicago, IL.
- Crane, J.L., D.E. Smorong, D.A. Pillard, and D.D. MacDonald. 2002. Sediment remediation scoping project in Minnesota Slip, Duluth Harbor. EPA-905-R97-020. U.S. Environmental Protection Agency, Great Lakes National Program Office, Chicago, IL.
- Dahl, T.E. 1990. Wetlands losses in the United States 1780s to 1980s. U.S. Fish and Wildlife Service, Washington, DC
- Davis, T.E., G.J. Niemi, J. Kotar, and P.B. Hofslund. 1978. Assessment of habitat types and bird populations in the lower St. Louis River, Phase II. Report to the Metropolitan Interstate Committee, Duluth, MN.
- DeVore, Philip W. 1978. Fishery resources of the Superior-Duluth estuary. Center for Lake Superior Environmental Studies, University of Wisconsin, Superior.
- Duluth Seaway Port Authority. 2002. Port Facts. Accessed May 2002 from World Wide Web site <http://www.duluthport.com/seawayfactsus.html>.
- Eggers, S. D., and D. M. Reed. 1997. Wetland plants and plant communities of Minnesota and Wisconsin. U.S. Army Corps of Engineers, St. Paul MN.
- Epstein, Eric J., Emmet J. Judziewicz, and William A. Smith. 1997. Priority wetland sites of Wisconsin's Lake Superior Basin: An excerpt from Wisconsin's Lake Superior Coastal Wetlands Evaluation. Report to the Great Lakes National Program Office of the U.S. Environmental Protection Agency. Wisconsin's Natural Heritage Inventory Program, Bureau of Endangered Resources, Department of Natural Resources, Madison, WI.
- Farrand, W. R., and C.W. Drexler. 1985. Late Wisconsinan and Holocene history of the Lake Superior Basin. Pages 17-32 in *Quaternary Evolution of the Great Lakes*, P.F. Karrow and P.E. Calkin, eds. Geological Association of Canada Special Paper 30.
- Finley, Robert W. 1976. Original vegetation cover of Wisconsin. Cartographic Laboratory, University of Wisconsin, Madison, WI.
- Frelich, L.E. 1999. Range of natural variability in forest structure for the Northern Superior Uplands. Report to the Minnesota Forest Resources Council. Department of Forest Resources, University of Minnesota, Minneapolis MN.
- Frelich, L.E., and C.G. Lorimer. 1991. Natural disturbance regimes in hemlock-hardwood forests of the Upper Great Lakes region. *Ecological Monographs* 61(2):145-164.

Lower St. Louis River Habitat Plan

- Fritzen, John. 1978. The history of Fond du Lac and Jay Cooke State Park. St. Louis County Historical Society, Duluth, MN.
- Galatowitsch, S.M., N.O. Anderson, and P.D. Ascher. 1999. Invasiveness in wetland plants in temperate North America. *Wetlands* 19(4):733-755.
- Gibbs, J.P., F.A. Reid, and S.M. Melvin. 1992. Least bittern (*Ixobrychus exilis*). In *The Birds of North America*, A. Poole, P. Stettenheim, and F. Gill, eds. (No. 17.) Academy of Natural Sciences, Philadelphia, and American Ornithologists' Union, Washington, DC
- Hale, Cindy M., Lee E. Frelich, and Peter B. Reich. 2002. Impacts of invading European earthworms on understory plant communities in previously worm-free hardwood forests of Minnesota. In preparation.
- Haufler, J.B., C.A. Mehl, and G.J. Roloff. 1996. Using a coarse-filter approach with species assessment for ecosystem management. *Wildlife Society Bulletin* 24(2):200-208.
- Hawrot, R.Y., and F.J. Nicoletti. 1999. Migratory and breeding bird population assessment document: Minnesota Point Protection M-22. In *Minnesota Point environmental management plan*, Park Point Community Club, Duluth, MN.
- Heinselman, M.L. 1973. Fire in the virgin forests of the Boundary Waters Canoe Area, Minnesota. *Quaternary Research* 3:329-382.
- Henson, F.G., and R. Newman. 2000. Effect of temperature on growth at ration and gastric evacuation rate of ruffe. *Trans. Am. Fish. Soc.* 129:552-560.
- Herkert, J. R., ed. 1992. Endangered and threatened species of Illinois: Status and distribution. Vol. 2: Animals. Illinois Endangered Species Protection Board, Springfield, IL.
- Howarth, F.G., and G.W. Ramsey. 1991. The conservation of island insects and their habitat. Pages 71-107 in *Conservation of insects and their habitats*, N.M. Collins and J.A. Thomas, eds. London: Academy Press.
- IJC (International Joint Commission). 1989. Great Lakes Water Quality Agreement of 1978 (as amended by Protocol signed November 18, 1987). IJC, Windsor, ON.
- IT Corp. (International Technology Corporation). 1997. Remedial investigation data report, sediment operable unit, St. Louis River/Interlake/Duluth Tar site. Volume 1. Prepared for the Interlake Corporation, Lisle, IL, by IT Corp., St. Paul, MN.
- Kellner D K., T. Kroska, and K. Plass. 1999. Historic reconstruction of property ownership and land uses along the Lower St. Louis River. St. Louis River Citizens Action Committee, Duluth, MN.

Lower St. Louis River Habitat Plan

- Kelly, J.R., J.A. Morrice, A.M. Cotter, M.L. Knuth, A.S. Trebitz, and R.L. Anderson. 1999. A new type of "lake effect"? Seiche-driven nutrient input to a coastal Great Lake ecosystem. ASLO Aquatic Sciences Meeting, Santa Fe, NM, February 1-5, 1999.
- King, P. 1999. Lake Superior/Duluth-Superior Harbor toxics loading study. Environmental Outcomes Division, Minnesota Pollution Control Agency, St. Paul, MN.
- Kraft, C. 1993. Early detection of the zebra mussel (*Dreissena polymorpha*). Pages 705-714 in Zebra mussels: Biology, impacts, and control, T.F. Nalepa and D.W. Schloesser, eds. Boca Raton, FL: CRC Press, Inc.
- Kruger, S., and R. Mishaga. 1996. A combined coarse-filter, fine-filter approach to implementing ecosystem management at the U.S. Army's Yakima Training Center in central Washington. Bulletin of the Ecological Society of America 77 (3 SUPPL. PART 2):245. Annual Combined Meeting of the Ecological Society of America on Ecologists/Biologists as Problem Solvers, Providence, Rhode Island, USA, August 10-14, 1996.
- Landmesser, C. W., and T.C. Johnson. 1982. Seismic reflection study of recessional moraines beneath Lake Superior and their relationship to regional deglaciation. Quaternary Research 17:173-190.
- Lindgren, J.R., P.W. Ongstad, and J.R. Spurrier. 1997. The St. Louis bay fishery, 1986-1994. Minnesota Department of Natural Resources, Division of Fisheries, Study 4 Report, Job 422, St. Paul, MN.
- Lodge, D.M., and J.G. Lorman. 1987. Reductions in submerged macrophyte biomass and species richness by the crayfish *Orconectes rusticus*. Can. J. Fish. Aquat. Sci. 44:591-597.
- Lusignan, Paul R. 1983. Superior intensive survey report. City of Superior Community Development Office, Superior, WI.
- Marschner, Francis J. 1974. The original vegetation of Minnesota. North Central Forest Experiment Station, Forest Service, U.S. Dept. of Agriculture, St. Paul, MN.
- Mayo, K.R., J.H. Selgeby, and M.E. McDonald. 1998. A bioenergetics modeling evaluation of top-down control of ruffe in the St. Louis River, western Lake Superior. J. Great Lakes Res. 24(2):329-342.
- MDH (Minnesota Department of Health). 2002. Minnesota fish consumption advisory. Health Risk Assessment Unit, Minnesota Department of Health, St. Paul, MN.
- MIC (Metropolitan Interstate Committee). 1985. Superior-Duluth Harbor natural resources management program. Arrowhead Regional Development Commission, Duluth, MN.
- Mills, E.L., J.H. Leach, J.T. Carlton, and C.L. Secor. 1993. Exotic species in the Great Lakes: A history of biotic crises and anthropogenic introductions. J. Great Lakes Res. 19:1-54.

Lower St. Louis River Habitat Plan

- MPCA and WDNR. 1992. The St. Louis River System remedial action plan: Stage One. Minnesota Pollution Control Agency, St. Paul, MN, and Wisconsin Department of Natural Resources, Madison, WI.
- MPCA and WDNR. 1995. The St. Louis River System remedial action plan: Stage Two. Minnesota Pollution Control Agency, St. Paul, MN, and Wisconsin Department of Natural Resources, Madison, WI.
- NatureServe. 2001. International classification of ecological communities: Terrestrial vegetation. Natural Heritage Central Databases. Arlington, VA: NatureServe.
- NatureServe Explorer. 1999. An online encyclopedia of life. Version 1.6. Accessed July 1999 from World Wide Web site <http://www.natureserve.org/explorer>.
- Nichols, S.J., J. French III, G. Kennedy, G. Black, J. Allen, E. Crawford, M. Blouin, J. Hickey, S. Chernyak, R. Haas, and M. Thomas. 2002. Threats to recovery of lake sturgeon (*Acipenser fulvescens*) in the lower St. Clair River. *J. Great Lakes Res.* (Accepted - In press).
- Niemi, G.J., T.E. Davis, and P.B. Hofslund. 1979. Distribution of habitats and birds in the St. Louis River Estuary, Minnesota and Wisconsin. Report to the U. S. Fish and Wildlife Service, St. Paul, MN.
- Niemi, G.J., J. Solin, D. Watters, and P. Wolter. 2000. A report on breeding bird inventory of the St. Louis River, Minnesota and Wisconsin, 1999, to The Nature Conservancy. Natural Resources Research Institute, University of Minnesota-Duluth, Duluth, MN.
- Noss, R.F., and A.Y. Cooperrider. 1994. Saving nature's legacy: Protecting and restoring biodiversity. Washington, DC: Island Press.
- NRCS (Natural Resources Conservation Service). 1988. Erosion and sedimentation in the Nemadji River Basin: Nemadji River Basin final report. U.S. Forest Service, Natural Resources Conservation Service, Duluth, MN.
- Nute, Grace Lee. 1944. Lake Superior, a volume in the American Lakes Series. Indianapolis and New York: The Bobbs-Merrill Company Publishers.
- Ojakangas, R. W., and C.L. Matsch. 1982. Minnesota's geology. Minneapolis, MN: University of Minnesota Press.
- Olsen, T.M., D.M. Lodge, G.M. Capelli, and R.J. Houlihan. 1991. Mechanisms of impact of an introduced crayfish (*Orconectes rusticus*) on littoral congeners, snails, and macrophytes. *Can. J. Fish. Aquat. Sci.* 48(10):1853-1861.
- Panzer, R., and M.W. Schwartz. 1998. Effectiveness of a vegetation-based approach to insect conservation. *Conservation-Biology* 12(3):693-702.
-

Lower St. Louis River Habitat Plan

- Park Point Community Club. 1999. Minnesota Point environmental management plan. Park Point Community Club, Duluth, MN.
- Pearson, S. 1999. St. Louis River Estuary colonial bird program 1999. Prepared for MN DNR. Department of Natural Resources, Brule, WI.
- Penning, W.L., and F.J. Cuthbert. 1993. The history of colonial waterbird management in the Duluth-Superior Harbor 1937-1990. *Loon* 65(4):163-174.
- Persaud, D., R. Jaagumagi, and A. Hayton. 1993. Guidelines for the protection and management of aquatic sediment quality in Ontario. Standards Development Branch, Ontario Ministry of Environment and Energy, Toronto, ON.
- Peterjohn, B.G., and J.R. Sauer. 1997. Population trends of black terns from the North American Breeding Bird Survey, 1966-1996. *Colonial-Waterbirds* 20(3):566-573.
- Phillips, G.L., W.D. Schmid, and J.C. Underhill. 1982. Fishes of the Minnesota region. Minneapolis, MN: University of Minnesota Press.
- Prater, B., and M. Anderson. 1977. A 96-hour sediment bioassay of Duluth and Superior Harbor basins (Minnesota) using *Hexagenia limbata*, *Asellus communis*, *Daphnia magna*, and *Pimephales promelas* as test organisms. *Bull. Environ. Contam. Toxicol.* 18:159-169.
- Redman, S., and T. Janisch. 1995. Newton Creek system sediment contamination site characterization report. Sediment Management and Remediation Techniques Program, Wisconsin Department of Natural Resources, Madison, WI.
- Ricciardi, A., and H. MacIsaac, 2000. Recent mass invasion of the North American Great Lakes by Ponto-Caspian species. *Trends in Ecology and Evolution* 15:62-65.
- Russell, R. 1983. The piping plover in the Great Lakes region. *American Birds* 37:951-955.
- Saltonstall, K. 2002. Cryptic invasion by a non-native genotype of the common reed, *Phragmites australis*, into North America. *Proceedings of the National Academy of Sciences of the United States of America*. [print] February 19, 2002; 99(4):2445-2449.
- Sauer, J.R., and S. Droege. 1992. Geographical patterns in population trends of neotropical migrants in North America. Pages 26-42 in *Ecology and Conservation of Neotropical Migrant Landbirds*, J.M. Hagan, III and D.W. Johnston, eds. Washington, DC: Smithsonian Institution Press.
- Sauer, J.R., J.E. Hines, I. Thomas, J. Fallon, and G. Gough. 2000. The North American breeding bird survey, results and analysis 1966 - 1999. Version 98.1. USGS Patuxent Wildlife Research Center, Laurel, MD. Accessed January 2002 from World Wide Web site <http://www.mbr-pwrc.usgs.gov/bbs/bbs99.html>.

Lower St. Louis River Habitat Plan

- Sauer, J.R., J.E. Hines, and J. Fallon. 2001. The North American breeding bird survey, results and analysis 1966 - 2000. Version 2001.2. USGS Patuxent Wildlife Research Center, Laurel, MD. Accessed January 2002 from World Wide Web site <http://www.mbr-pwrc.usgs.gov/bbs/bbs.html>.
- Schoolcraft, Henry R. 1855. Summary narrative of an exploratory expedition to the source of the Mississippi River in 1820, resumed and completed by the discovery of its origin in Itasca Lake in 1832. Philadelphia: Lippincott, Grambo & Co.
- Schram, S.T., J.R. Lindgren, and L.M. Evrard. 1999. Reintroduction of lake sturgeon in the St. Louis River, western Lake Superior. *North Am. J. Fisheries Management* 19:815-823.
- Schubauer-Berigan, M., and J.L. Crane. 1996. Preliminary contaminant assessment of the Thomson, Forbay, and Fond du Lac reservoirs. Water Quality Division, Minnesota Pollution Control Agency, St. Paul, MN.
- Schubauer-Berigan, M., and J.L. Crane. 1997. Survey of sediment quality in the Duluth/Superior Harbor: 1993 sampling results. EPA 905-R97-005. U.S. Environmental Protection Agency, Great Lakes National Program Office, Chicago, IL.
- Schueler, T. 1987. Controlling urban runoff: a practical manual for planning and designing urban best management practices. Metropolitan Washington Council of Governments, Washington DC.
- Schuldt, J.A., C. Richards, J. Gangl, and R.M. Newman. In prep. Effects of Eurasian ruffe on fish food resources in mesocosms.
- SEH (Short Elliot Hendrickson). 1994. Site investigation report and closure plan, prepared for Fraser Shipyards, Inc., Superior WI, by SEH, Chippewa Falls, WI.
- Selgeby, J. 1998. Predation by ruffe (*Gymnocephalus cernuus*) on fish eggs in Lake Superior. *J. Great Lakes. Res.* 24(2):304-308.
- Snetsinger, S.D. 2000. Land cover change in the Great Lakes region: from pre-European settlement to present. M.S. Thesis, University of Wisconsin, Madison. 113 pp.
- Snetsinger, S., and S. Ventura. 2000. Land cover change in the Great Lakes region from mid-nineteenth century to present. Great Lakes Ecological Assessment. Accessed January 2002 from World Wide Web site <http://www.ncrs.fs.fed.us/gla/>.
- Stearns, F.W. 1949. Ninety years of change in a northern hardwood forest in Wisconsin. *Ecology* 30(3):350-358.
- TNC (The Nature Conservancy). 2000. The 5-S framework for site conservation: A practitioner's handbook for site conservation planning and measuring conservation success. Arlington, VA: The Nature Conservancy.

Lower St. Louis River Habitat Plan

- U.S. ACOE (Army Corps of Engineers). 1987. Great Lakes water level facts. Detroit District, U.S. Army Corps of Engineers, Detroit, MI.
- U.S. ACOE (Army Corps of Engineers). 1997. Phase II report. Draft dredged material management plan study and environmental impact statement. Detroit District, U.S. Army Corps of Engineers, Detroit, MI.
- U.S. EPA (Environmental Protection Agency). 1983. Final report: Results of the nationwide urban runoff project. U.S. Environmental Protection Agency, Washington DC.
- U.S. FWS (Fish and Wildlife Service). 2001. Endangered and threatened wildlife and plants: Final determination of critical habitat for the Great Lakes breeding population of the piping plover. Federal Register 66, no. 88, (7 May 2001): 22938-22969.
- Verry E. S., J. S. Lewis, and K. N. Brooks. 1983. Aspen clearcutting increases snowmelt and storm flow peaks in north central Minnesota. *Water Resources Bulletin* 19(1):59-67.
- Verry, E. S. 1986. Forest harvesting and water: the Lake States experience. *Water Resources Bulletin* 22(6):1039-1047.
- Walker, D.A., and S.P. Hall. 1976. Duluth-Superior Harbor cultural resources study. Archeology Department, Minnesota Historical Society, St. Paul, MN.
- WDNR (Wisconsin Department of Natural Resources). 2002. Hook into healthy fish. PUB-FH-505. Wisconsin Department of Natural Resources, Madison, WI.
- Whitney, G.G. 1986. Relation of Michigan's presettlement pine forests to substrate and disturbance history. *Ecology* 67(6):1548-1559.
- Zhuikov, M. 1999. Great Salt Lake Superior? Global warming's impacts predicted. *Seiche* newsletter, April. University of Minnesota, Minnesota Sea Grant Program, Duluth, MN.

APPENDICES

Appendix 1. List of Participants and Reviewers

Appendix 2. List of Plant Communities Mapped in the Lower St. Louis River

Appendix 3. List and Descriptions of Aquatic Habitat Types of the Lower St. Louis River and Their Ecological Values

Appendix 4. Conservation Target-Related Recommendations from St. Louis River Remedial Action Plan Stage Two

Appendix 5. List of Fish Species Native to the Lower St. Louis River and Their Requirements

Appendix 6. Fish Use of Aquatic Habitats in the Lower St. Louis River

Appendix 7. Birds of the Lower St. Louis River

Appendix 8. Summary of Threats Identified for Each Conservation Target in the Lower St. Louis River

Lower St. Louis River Habitat Plan

MAPS

Map 1. Locational Map of Project Area Boundary

Map 2. Presettlement Vegetation of the Lower St. Louis River Project Area

Map 3. Current Vegetation of the Lower St. Louis River Project Area

Map 4. St.Louis River Plant Communities and Aquatic Habitats

Map 5. Aquatic Habitat Types